

THE TREMFYA® FACT SHEET

Here are some facts about TREMFYA® that you can discuss with your dermatologist. He or she can answer any questions you have about the treatments used in patients with moderate to severe plaque psoriasis who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet light or UV light).

UNCOVER CLEARER SKIN THAT CAN LAST

In clinical studies at 16 weeks, at least 7 out of 10 patients saw at least 90% clearer skin, and more than 8 out of 10 patients were rated clear or almost clear. In a study, nearly 9 out of 10 people who saw at least 90% clearer skin at 28 weeks continued to see at least 90% clearer skin at 48 weeks while on TREMFYA®.

UNCOVER RESULTS AT 5 YEARS

In a study, nearly 7 out of 10 patients with 90% clearer skin at 16 weeks were still clearer at 5 years. At 1 year and thereafter, patients and healthcare providers knew that TREMFYA® was being used. This may have increased results. Results may vary.

UNCOVER SUPERIOR SKIN CLEARANCE

In clinical studies, TREMFYA® worked better than Humira® (adalimumab) at providing significantly clearer skin.

Humira® is a registered trademark of AbbVie Inc.

UNCOVER COMPLETELY CLEAR SKIN

In another clinical study, 5 out of 10 patients saw completely clear skin.

UNCOVER CLEARER SKIN

RESULTS FROM CLINICAL STUDIES AT WEEK 16

≥90% CLEARER SKIN



CLEAR OR ALMOST CLEAR



Individual results may vary. TREMFYA® is not for everyone. Only you and your dermatologist can decide if TREMFYA® is right for you.

TREMFYA® is a prescription medicine used to treat adults with moderate to severe plaque psoriasis who may benefit from taking injections or pills (systemic therapy) or phototherapy (treatment using ultraviolet light or UV light).

TREMFYA® TREATMENT EVERY 8 WEEKS, AFTER 2 STARTER DOSES:

TREMFYA® is given as a 100 mg injection under your skin that you only need to take every 8 weeks, after 2 starter doses at weeks 0 and 4. That's about 2 whole months that you don't have to think about your treatment schedule.

SELECTED IMPORTANT SAFETY INFORMATION

TREMFYA® is not for everyone; only your doctor can decide if it's right for you. Do not use if you are allergic to TREMFYA®. TREMFYA® is a prescription medicine that may cause serious side effects, including serious allergic reactions and infections. TREMFYA® affects your immune system. It may increase your risk of infections and lower your ability to fight them. Please read the Important Safety Information on the next page and the [Medication Guide](#) for TREMFYA® to learn more about these and other risks for TREMFYA®. Discuss any questions you have with your doctor.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about TREMFYA®? TREMFYA® is a prescription medicine that may cause serious side effects, including:

- **Serious Allergic Reactions.** Stop using TREMFYA® and get emergency medical help right away if you develop any of the following symptoms of a serious allergic reaction:
 - fainting, dizziness, feeling lightheaded (low blood pressure)
 - swelling of your face, eyelids, lips, mouth, tongue or throat
 - trouble breathing or throat tightness
 - chest tightness
 - skin rash, hives
 - itching
- **Infections.** TREMFYA® may lower the ability of your immune system to fight infections and may increase your risk of infections. Your healthcare provider should check you for infections and tuberculosis (TB) before starting treatment with TREMFYA® and may treat you for TB before you begin treatment with TREMFYA® if you have a history of TB or have active TB. Your healthcare provider should watch you closely for signs and symptoms of TB during and after treatment with TREMFYA®.

Tell your healthcare provider right away if you have an infection or have symptoms of an infection, including:

- fever, sweats, or chills
- muscle aches
- weight loss
- cough
- warm, red, or painful skin or sores on your body different from your psoriasis
- diarrhea or stomach pain
- shortness of breath
- blood in your phlegm (mucus)
- burning when you urinate or urinating more often than normal

Do not take TREMFYA® if you have had a serious allergic reaction to guselkumab or any of the ingredients in TREMFYA®.

Before using TREMFYA®, tell your healthcare provider about all of your medical conditions, including if you:

- have any of the conditions or symptoms listed in the section **“What is the most important information I should know about TREMFYA®?”**
- have an infection that does not go away or that keeps coming back.
- have TB or have been in close contact with someone with TB.
- have recently received or are scheduled to receive an immunization (vaccine). You should avoid receiving live vaccines during treatment with TREMFYA®.
- are pregnant or plan to become pregnant. It is not known if TREMFYA® can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if TREMFYA® passes into your breast milk.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

What are the possible side effects of TREMFYA®?

TREMFYA® may cause serious side effects. See “What is the most important information I should know about TREMFYA®?”

The most common side effects of TREMFYA® include: upper respiratory infections, headache, injection site reactions, joint pain (arthralgia), diarrhea, stomach flu (gastroenteritis), fungal skin infections, herpes simplex infections, and bronchitis.

These are not all the possible side effects of TREMFYA®. Call your doctor for medical advice about side effects.

Use TREMFYA® exactly as your healthcare provider tells you to use it.

Please read the full [Prescribing Information](#), including [Medication Guide for TREMFYA®](#), and discuss any questions that you have with your doctor.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch, or call 1-800-FDA-1088.

cp-82626v3